

ARL10B

PDB:1ZD9

Revision

Revision Type:created

Revised by:created

Revision Date:created

Entry Clone Accession:AT20-E2

Entry Clone Source:MGC

SGC Clone Accession:HPC003-F10

Tag:mgsshhhhhhsglvp*r*gs

Host:E. coli BL21 (DE3)

Construct

Prelude:

Sequence:

mgsshhhhhhsglvp*gsKEEMELTVGLQYSGKTFVNVIASGQFNEDMIPTVGFNMRKITKGNVTIKLWDIGGQPRFRSMWERY
CRGVSAIVYMVAADQEKEASKNELHNLLDKPQLQGIPVLVLGNKRDLPGALDEKELIEKMNLSAIQDREICCYSCKEKDNDI
TLQWLIQHSKSRRS

Vector:p28a-thrombin-lic

Growth

Medium:

Antibiotics:

Procedure:We prepared a seed culture by inoculating freshly transformed E. coli cells (BL21 DE3) into 80 mL of Luria-Bertani medium. After overnight, the seed culture was inoculated into 1.8 L of Terrific Broth medium in the presence of 50 µg/ml of kanamycin at 37°C and grown to an OD600 of 4.63. Cells were then induced by isopropyl-1-thio-D-galactopyranoside at the final concentration of 1.5 mM and grown overnight at 20°C in a LEX bubbling system.

Purification

Procedure

The supernatant was passed through DE52 (Whatman) column equilibrated with the binding buffer and then loaded onto 3 ml Ni-NTA column (Qiagen) equilibrated with the same binding buffer at 4oC. The Ni-NTA column was washed with 150 ml of the wash buffer (10mM Tris pH7.5, 0.5 M NaCl, 5% glycerol, 30 mM imidazole) and the protein was eluted with 15 ml of the elution buffer (10mM Tris pH7.5, 0.5 M NaCl, 5% glycerol, 250 mM imidazole). The eluate was dialyzed overnight against a buffer containing 10 mM Tris pH7.5, 0.5 M NaCl, 5% glycerol. The protein concentration was estimated based on the extinction coefficient of the protein, 22430 at

280 nm. Five molar equivalents of GDP, 5 mM TCEP and 5 mM MgCl₂ were added to the purified protein before concentration. The protein was concentrated using an Amicon Ultra centrifugal filter to the final volume of 1 ml and the concentration of 30 mg/ml. About 55 mg of protein was obtained from 1.8 L of cell culture.

Extraction

Procedure

Cultures were centrifuged and the cell pellets were suspended in 100 ml of the binding buffer (10 mM Tris pH 7.5, 0.5 M NaCl, 5% glycerol, 5 mM imidazole) with a protease inhibitor cocktail (0.1 mM M benzamidine-HCl and 0.1 mM phenylmethyl sulfonyl fluoride) and flash frozen. The thawed cell pellet was lysed by a combination of 0.5% CHAPS (Sigma) and sonication. The lysate was centrifuged at 15000 rpm for 30 min and the supernatant was used for subsequent steps of purification.

Concentration: 30 mg/mL

Ligand

MassSpec:

Crystallization: Purified His-tagged ARL10B was crystallized using the sitting drop vapor diffusion method. Crystals grew in one day when the protein (30 mg/mL) was mixed with the reservoir solution in a 1:1 volume ratio, and the drop was equilibrated against a reservoir solution containing 30 % PEG 4000, 0.1 M Tris, pH 8.5 and 0.2 M LiSO₄.

NMR Spectroscopy:

Data Collection:

Data Processing: