

Cryptosporidium parvum hypothetical protein cgd2_2020

PDB:2PD0

Revision

Revision Type:created

Revised by:created

Revision Date:created

Entry Clone Accession:cgd2_2020

Entry Clone Source:Cryptosporidium parvum strain Iowa genomic DNA

SGC Clone Accession:cp-PFA0230c, plate MS:C1

Tag:mgsshhhhhhsglvpr(*)gs

Host:E. coli BL21-(DE3)-R3-pRARE2

Construct

Prelude:

Sequence:

gsKLLGVNSFALRQFVEGYRGSYIPRMSPYFLRNVNYYIIENNPTLVGYADFCKHIFIPNFTAKQSIVKITNEKYIKTGYIS
RRDEEIPVLSRWFPKDSPPASQLIKSKYLDIILYSKEQCEKESSIMNCCLQDILDDREKNPDWYIISIKAQNESFEVPMEPITILRN
TLIEEGGSGVPLKREKYLESVFWKEHAISS

Vector:pET28A-lic

Growth

Medium:M9 Selmet kit

Antibiotics:100 microG/mL ampicillin and 34 microG/mL chloramphenicol

Procedure:A single colony was inoculated into 10 mL of LB with of Antibiotics and incubated with shaking at 250 rpm overnight at 37 degC. The culture was transferred into 50 mL of TB with Antibiotics in a 250 mL shaking flask and incubated at 37 degC for 3 hours. The culture was then transferred into 1.8 L of above-specified growth medium with Antibiotics and 0.3 mL of antifoam (Sigma) in a 2L bottle and cultured using the LEX system to an OD600 of ~1, cooled to 15 degC and induced with 0.5 mM isopropyl-1-thio-D-galactopyranoside (IPTG) overnight at 20 degC.

Purification

Procedure

The cleared lysate was loaded onto a column prepacked with 10 g DE52 (Whatman) anion exchange resin (previously activated with 2.5 M NaCl and equilibrated with Binding Buffer); and subsequently onto a 1.0-2.5 mL Ni-NTA (Qiagen) column pre-equilibrated with Binding Buffer at approximately 1-1.5 mL/min. The volume of the Ni-NTA resin was pre-determined by the predicted protein yield from test expression analysis. After the lysate was loaded, the DE52 was further washed with 20 mL of Binding Buffer. The Ni-NTA column was then washed with 200

mL of Wash Buffer at 2-2.5 mL/min. After washing, the protein was eluted with 15 mL of Elution Buffer. EDTA was immediately added to the elution fraction to 1 mM; and TCEP was added to 1-5 mM after approximately 15 more minutes.

The His6-tag was cleaved with thrombin overnight at 4 degC in the presence of 1 mM TCEP and dialysed into Crystal Buffer (10 mM HEPES, pH 7.5, 500 mM NaCl). The cleaved sample was loaded onto a Sephadex S75 26/60 column equilibrated with Gel Filtration Buffer. The fractions from the peak eluting at 164 mL corresponding to monomeric protein were pooled and concentrated using a 15 mL Amicon Ultra centrifugal filter device (Millipore). The protein sample identity and purity were evaluated by mass spectroscopy and SDS-PAGE gel. The concentrated protein was stored at 4 degC.

Extraction

Procedure

Cells were resuspended to approximately 40 mL/L of cell culture in Binding Buffer with protease inhibitor (1 mM benzamidine-HCl and 1 mM phenylmethyl sulfonyl fluoride, PMSF).

Resuspended pellets stored at -80 degC were thawed overnight at 4 degC on the day before purification. Prior to mechanical lysis, each pellet from 1 L of culture was pretreated with 0.5% CHAPS and 500 units of benzonase for 40 minutes at room temperature. Cells were mechanically lysed with a microfluidizer (Microfluidizer Processor, M-110EH) at approximately 18000 psi; and the cell lysate was centrifuged at ~75000 x g for 20 minutes at 10 degC.

Concentration:

Ligand

MassSpec:

Crystallization: The protein was crystallized at 20 degC in 21% P3350, 0.2 M LiCl, 0.1 M MES pH 5.5 using the sitting drop vapor diffusion method.

NMR Spectroscopy:

Data Collection:

Data Processing: