

# Tg-CyP16L

**PDB:**3BKP

## Revision

**Revision Type:**created

**Revised by:**created

**Revision Date:**created

**Entry Clone Accession:**49.m03261

**Entry Clone Source:**

**SGC Clone Accession:**49.m03261:S10-S223:B10

**Tag:**mhhhhhhsgrenlyfqg

**Host:**Ros-Ox

## Construct

**Prelude:**

**Sequence:**

STRGVVLHTSLGDLVELWARECPLACRNFVQLCLEGYYNTIFHRVVKDFIVQGGDPTGTGRGGADTTFDGKPFDVETHPRLKFR  
YRGLVGVANLGRSSKDAENDERGRSLGTNGNQFFITLARADVNNAYTLFGKVTGHTLYNLMKFNDLEVKGKEDRPMTPPFIKSVDVL  
WNPFEDLVPRRLPDAPPAQKDERKRARAAGGRTEGRARS

**Vector:**p15-mhl

## Growth

**Medium:**TB

**Antibiotics:**

**Procedure:**49.m03261 was expressed in E. coli BL21-(DE3)-Rosetta-Oxford cells in Terrific Broth (TB) in the presence of ampicillin/chloramphenicol (50 microg/mL and 25 microg/mL respectively). A single colony was inoculated into 10 mL of LB with of ampicillin/chloramphenicol (50 microg/mL and 25 microg/mL respectively) in a 50 mL Falcon tube and incubated with shaking at 250 rpm overnight at 37 degC. The culture was transferred into 50 mL of TB with 50 microg/mL ampicillin in a 250 mL shaking flask and incubated at 37 degC for 3 hours. Then the culture was transferred into 1.8 L of TB with 50 microg/mL kanamycin and 0.3 mL of antifoam (Sigma) in a 2 L bottle and cultured using the LEX system to an OD 600 of ~5, cooled to 15 degC, and induced with 0.5 mM isopropyl-1-thio-D-galactopyranoside (IPTG) overnight at 15 degC.

## Purification

**Procedure**

**STEP1:**The cleared lysate was loaded onto a column prepacked with 10 g DE52 (Whatman) anion exchange resin (previously activated with 2.5 M NaCl and equilibrated with Binding

Buffer); and subsequently onto a 3 mL Ni-NTA (Qiagen) column pre-equilibrated with Binding Buffer at approximately 1  $\text{mL}/\text{min}$ . The volume of the Ni-NTA resin was pre-determined by the predicted protein yield from test expression analysis. After the lysate was loaded, the DE52 was further washed with 20 mL of Binding Buffer. The Ni-NTA column was then washed with 200 mL of Wash Buffer at 2  $\text{mL}/\text{min}$ . After washing, the protein was eluted with 15 mL of Elution Buffer. EDTA was immediately added to the elution fraction to 1 mM; and TCEP was added to 0.5 mM after approximately 15 more minutes.

**STEP2:** The sample was loaded onto a Sephadex S75 26/60 gel filtration column pre-equilibrated with 10 mM HEPES, pH 7.5 and 500 mM NaCl. The collected fractions corresponding to the correct eluted protein peak were concentrated using a 15 mL Amicon Ultra centrifugal filter device (Millipore). The protein sample identity were evaluated by mass spectroscopy. The concentrated sample (47 mg/ml) was stored at 4 degC.

## **Extraction**

### **Procedure**

The culture was harvested by centrifugation. Pellets from 4 L of culture were resuspended to approximately 40 mL/L of cell culture in Binding Buffer with the addition of protease inhibitors (1 mM benzamidine and 1 mM phenylmethyl sulfonyl fluoride (PMSF)). Resuspended pellets stored at -80 degC were thawed overnight at 4 degC on the day before purification. Prior to mechanical lysis, each pellet from 1 L of culture was pretreated with 0.5 % CHAPS and 500 units of benzonase for 40 minutes at room temperature. Cells were mechanically lysed with a microfluidizer (Microfluidizer Processor, M-110EH) at approximately 18000 psi; and the cell lysate was centrifuged using a Beckman JA-25.50 rotor at ~75000 x g (24000 rpms) for 20 minutes at 10 degC.

### **Concentration:**

### **Ligand**

### **MassSpec:**

**Crystallization:** MAY63L-H:E1025% glycerol The protein was crystallized at 20 degC in 25% w/v PEG 3350 0.2 M Sodium Chloride 0.1M HEPES pH - 7.5 using the sitting drop vapor diffusion method.

### **NMR Spectroscopy:**

### **Data Collection:**

### **Data Processing:**