

MERTK

PDB:3BRB

Revision

Revision Type:created

Revised by:created

Revision Date:created

Entry Clone Accession:axl2.LIFESEQ1708542

Entry Clone Source:Open Biosystems

SGC Clone Accession:SDC037B11

Tag:N-terminal: MGSSHHHHHSSGLVPRGS

Host:BL21 (DE3)

Construct

Prelude:

Sequence:

MGSSHHHHHSSGLVPRGSEELQNKLEDVVIDRNLLILGKILGEGEFGSVMEGNLKQEDGTSLKAVKTMKLDNSSQREIEFLSEA
ACMKDFSHPNVIRLLGVCIE MSSQGIPKPMVILPFMKGDLHTYLLYSRLETGPKH IPLQTLKFMVDIALGMEYLSNRNFLHDLA
ARNCMLRDDMTVCVADFGLSKKIYSGDYYRQGRIAKMPVKWIAIESLADRVTTSKSDVAFGVTMWEIATRGMTPYPGVQNHEMYDY
LLHGHLKQPEDCLDELYEIMYSCWRTDPLDRPTFSVRLQLEKLLESLPDV

Vector:pET28a-LIC

Growth

Medium:TB with 50 μ g/mL kanamycin

Antibiotics:

Procedure:A small overnight culture containing 50 μ g/mL kanamycin was used to inoculate TB media containing the same concentration of antibiotics. Cultures were grown at 37°C for about 6 hours until the OD₆₀₀ reached ~0.8, the temperature was adjusted to 15°C, and expression was induced using 0.1mM IPTG overnight. Cells were harvested by centrifugation and frozen.

Purification

Procedure

Clarified supernatant was mixed with 5.0 ml 50% Talon resin slurry (Clonetech), rotated for 1 hour at 4 oC, and then loaded into a column. Ten column volumes of washing buffer were used for washing before elution with 7 mL elution buffer.

Gel-filtration was conducted using AKTAexpress (18-6645-05, GE Healthcare) with XK 16x65 columns (part numbers 18-1031-47 and 18-6488-01, GE Healthcare) packed with HighLoad Superdex 200 resin (10-1043-04, GE Healthcare). Pre-equilibration was done with gel filtration

buffer at a flow rate of 3 mL/min. 7 mL of protein sample was loaded onto the column at 1.5 mL/min, and 2 mL fractions are collected in 96-well plates (VWR 40002-012) using peak fractionation protocols with the following parameters: (Slope; min. peak width 0.833 min; level 0.000 mAU; peak start slope 10.000 AU/min; peak end slope 20.000 AU/min). Fractions were analyzed for purity using SDS-PAGE and those containing pure Mer/axl2 were pooled.

Purified protein was concentrated using 15 mL concentrators with a 10,000 molecular weight cut-off (Amicon Ultra-15, UFC900524, Millipore) at 3750 rpm, typically resulting in a final concentration of 7-10 mg/mL.

Extraction

Procedure

Frozen cell pellets obtained from 2L culture were thawed, resuspended in 50 mL lysis buffer, homogenized with Ultra-Turrax T8 homogenizer (IKA Works) at maximal setting for 30-60 seconds, and twice passed through a microfluidizer (Microfluidics M110EH) at 18,000 psi. The lysate was clarified by centrifugation (JA25.50 rotor, Avanti J-20 XPI, Beckman Coulter) for 20 minutes at 69,673 x g.

Concentration: 7-10 mg/mL

Ligand

MassSpec: Calculated MW: 35841.4 g/mol

Crystallization: Mer protein (38 mg/ml) was pre-incubated with 2.5 mM ATP and 10 mM MgCl₂ for 3h at room temperature.

Crystals were obtained at 14 oC using the hanging-drop vapour-diffusion method by mixing 2 ul Mer solution with 2 ul reservoir (100 mM Tris-HCl pH 8.5, 200 mM MgCl₂, 29% PEG 400).

Cryo buffer was composed of glycerol, ethylene glycol, glucose, and fructose.

NMR Spectroscopy:

Data Collection:

Data Processing: