

ATAD2

PDB:3DAI

Revision

Revision Type:created

Revised by:created

Revision Date:created

Entry Clone Accession:gi|24497618

Entry Clone Source:synthetic

SGC Clone Accession:

Tag:mhahhhhhssggvdlgtenlyfq*s(m) TEV-cleavable (*) N-terminal his6 tag.

Host:BL21(DE3)-R3-pRARE2

Construct

Prelude:

Sequence:

mhahhhhhssggvdlgtenlyfq*smQEEDTFRELRIFLRNVTHLRAIDKRFRVFTKPVDPDEVPDYVTVIKQPMDSLSSVISKIDLHKY
LTVKDYLRIDLICSNALEYNPDRDPGDRLLIRHRACALRDTAYAIKEELDEDFEQLCEEIQESR

Vector:pNIC28-Bsa4

Growth

Medium:

Antibiotics:

Procedure:The expression plasmid was transformed into the host strain and plated on LB-agar containing 50 µg/ml kanamycin and 35 µg/ml chloramphenicol. Several colonies were combined to inoculate a 1ml culture in TB (+ 50 µg/ml kanamycin, 35 µg/ml chloramphenicol). The culture was grown overnight, glycerol was added to 15% v/v (from a 60% stock), and the resulting glycerol stock was frozen at -80°C in 100 µl aliquots. When required a 50 ml overnight culture was started by inoculating a loopful of cells from the glycerol stock into LB medium containing antibiotic as above. 5ml this overnight culture was used to inoculate similar 1 litre cultures (4L total) which were grown at 37°C until the OD600 reached ~0.5 at which point the temperature was adjusted to 18°C. Expression was induced overnight using 0.5 mM IPTG. The cells were collected by centrifugation and the pellet resuspended in binding buffer and frozen. (Binding buffer: 50mM HEPES pH 7.5; 500 mM NaCl; 5 mM imidazole, 5% glycerol.)

Purification

Procedure

Column 1: Ion exchange - Nucleic acid removal. DEAE cellulose (DE52, Whatman), 10 g of resin in 2.5 x 20 cm column. The resin was hydrated in 2.5M NaCl, washed with 20 ml binding

buffer prior to loading the sample. Buffers: 50mM HEPES pH 7.5; 500 mM NaCl; 5 mM imidazole. Procedure: Supernatant was applied by gravity flow, followed by a wash with 50 ml binding buffer. The column flowColumn 2: Ni-affinity. Ni-NTA (Qiagen), 5 ml of 50% slurry in 1.5 x 10 cm column, washed with binding buffer. Buffers: Binding buffer: 50 mM HEPES pH 7.5, 500 mM NaCl, 5 mM imidazole, 5% glycerol; Wash buffer: 50 mM HEPES pH 7.5, 500 mM NaCl, 30 mM Imidazole, 5% glycerol; Elution buffer: 50 mM HEPES pH 7.5, 500 mM NaCl, 50 to 250 mM Imidazole, 5% Glycerol. Procedure: The flowthrough from column 1 was loaded by gravity flow on the Ni-NTA column. The column was then washed with 100 ml wash buffer at gravity flow. The protein was eluted by gravity flow by applying 5-ml portions of elution buffer with increasing concentration of imidazole (50 mM, 100 mM, 150 mM and 250 mM); fractions were collected until essentially all protein was eluted. 10 mM DTT was added for overnight storage together with TEV protease for cleavage of the N-terminal hexahistidine tag. Column 3: Size Exclusion Chromatography. Superdex S200 16/60 HiLoad Buffers: 25 mM HEPES, pH 7.5; 150 mM NaCl, 0.5 mM TCEP. Procedure: Protein was applied to a S200 gel filtration column using an ÄKTAXpress system at 1.0 ml/min. Eluted proteins were collected in 1.8-ml fractions and analyzed on SDS-PAGE. 10 mM DTT was added for storage. Column 4: Ni-affinity. Ni-NTA (Qiagen), 2 ml of 50% slurry in 1.5 x 10 cm column, washed with binding buffer. Procedure: The elution from column 3 (TEV cleaved) was loaded by gravity flow on a Ni-NTA column to remove Ni-binding contaminants. The column was washed with 10 ml wash buffer (as column 2) and proteins eluted as column 2. 10 mM DTT was added for overnight storage.

Extraction

Procedure

Frozen pellets were thawed and fresh 1 mM PMSF and 0.5 mM TCEP were added. Cells were lysed by sonication. The lysate was centrifuged at 21,000 rpm for 60 minutes and the supernatant collected for purification.

Concentration: The cleaved purified protein was concentrated in a VivaSpin500 (5 K MWCO) to 12.2 mg/ml and stored at 4°C. The protein concentration was determined spectrophotometrically.

Ligand

MassSpec: Observed mass without histidine tag, 15430.6 (calculated mass without histidine tag, 15431).

Crystallization: Crystals were obtained using the vapor diffusion method. The protein was concentrated in gel filtration buffer to a protein concentration of 12.2 mg/ml. Sitting drops comprising 75 nl of the concentrated protein mixed with 75 nl of a well solution (2.00M (NH₄)₂SO₄; 0.1M BisTris pH 5.5) were equilibrated against well solution at 4°C. Crystals appeared within 2-3 days.

NMR Spectroscopy:

Data Collection: Crystals were cryo-protected using the well solution supplemented with 2M Li₂SO₄ and flash frozen in liquid nitrogen. Diffraction data were collected from a single crystal on a Rigaku FR-E SuperBright at a single wavelength of 1.5 Å. The structure was solved by molecular replacement and refined to 1.89 Å.

Data Processing: