

BRPF1

PDB:3L42

Revision

Revision Type:created

Revised by:created

Revision Date:created

Entry Clone Accession:GI:51173720

Entry Clone Source:MGC

SGC Clone Accession:

Tag:N-terminal: His-tag with integrated TEV protease site: MHHHHHHSSGRENLYFQG

Host:*E. coli* BL21 (DE3) codon plus RIL (Stratagene).

Construct

Prelude:

Sequence:

gEDSPLDALDLWAKCRGYPSPALIIDPKMPREGMFHHGVPPIPVPPLLEVKLGEQMTQEAREHLYVLFFDNKRTWQWLPRTKLVP
LGVNQQLDKEKMLEGRKSNIRKSVQIAYHRALQHRSKVQGEQS

Vector:pET28-MHL

Growth

Medium:

Antibiotics:

Procedure:BRPF1 was expressed in *E.coli* BL21 (DE3) codon plus in Terrific Broth medium in the presence of 50 µg/ml of kanamycin. Cell were grown at 37 degC to an OD600 of 1.5 and induced by isopropyl-1-thio-D-galactopyranoside (IPTG), final concentration 1 mM and incubated overnight at 15 degC.

Purification

Procedure

The lysate was loaded onto 5 ml HiTrap column (Amersham Biosciences), charged with Ni2+. The column was washed with 10 CV of Wash buffer, and the protein was eluted with elution buffer. The protein was dialyzed against buffer containing 20 mM HEPES, pH 7.4, 500 mM NaCl and 5% glycerol. TEV protease was added to combined fractions containing BRPF1. The protein was further purified to homogeneity by ion-exchange chromatography on Source 30S column (10x10) (Amersham Biosciences), equilibrated with buffer 20 mM PIPES, pH 6.5, and eluted with linear gradient of NaCl up to 500 mM concentration (20CV). Purification yield was 12 mg of the protein per 1L of culture.

Extraction

Procedure

Cells were harvested by centrifugation at 7,000 rpm. The cell pellets were frozen in liquid nitrogen and stored at -80 degC. For the purification the cell paste was thawed and resuspended in lysis buffer with protease inhibitor (0.1mM phenylmethyl sulfonyl fluoride, PMSF). The cells were lysed by passing through Microfluidizer (Microfluidics Corp.) at 20,000 psi.

Concentration:33.2 mg/ml.

Ligand

MassSpec:Expected MW is 15052.51 Da, measured mass is 15052.9917 Da.

Crystallization:Purified BRPF1 was crystallized using hanging drop vapor diffusion method at 20 °C by mixing 1 μ l of the protein solution (10 mg/mL) with 1 μ l of the reservoir solution containing 3.5 M NaForm, 0.1 M Tris-HCl, pH 8.5.

NMR Spectroscopy:

Data Collection:

Data Processing: