

MAPK

PDB:3N9X

Revision

Revision Type:created

Revised by:created

Revision Date:created

Entry Clone Accession:PB000659.00.0

Entry Clone Source:

SGC Clone Accession:PBG-PF11_0147:I110-S523:C2

Tag:mhhhhhhsgrenlyfqq

Host:BL21(DE3)-V2R-pACYC LamP

Construct

Prelude:

Sequence:

IKNVHPDNYIICKHLIGRGSGYVYLAYDKNTEKNVAIKKVNRMFEDLIDCKRILREITILNRLKSDYIIRLYDLIIPDDLKFDL
YIVLEIADSDLKKLFKTPIFLTEEHIKTILYNLLGENFIHESGIIHRLK PANCLLNQDCSVK CDFGLARTINSEKDTNIVNDLE
ENEERPGPHNKNLKKQLTSHVVTRWYRAPELILLQENYTKSIDIWSTGCIFAELLNMLQSHINDPTNRFPLFPGSSCFPLSPDRNSKK
VHEKSNRDQLNIIFNIIGTPTEDDLKNINKPEVIKYIKLFPHRKPINLKQKYPISDDGINLLESMLKFNPNKRTIDQALDHPYLK
DVRKKKLENFSTKKIILPFDDWMVLSETQLRYIFLKEVQSFHPELVIPSFTIHENNFTYNNNEKPSS

Vector:p15-mhl

Growth

Medium:TB

Antibiotics:

Procedure:Plasmodium berghei PBG-PF11_0147 was expressed in E. coli BL21(λDE3) V2R pACYC LamP in TB growth media in the presence of carbenicillin/chloramphenicol (100 microgram/mL and 34 microgram/mL, respectively). A single colony was inoculated into 25 mL of LB with of carbenicillin/chloramphenicol (100 microgram/mL and 34 microgram/mL respectively) in a 50 mL Falcon tube and incubated with shaking at 250 rpm overnight at 37degC. Then the culture was transferred into 900 mls of TB with 100 microgram/mL Carbenicillin and 34 microgram/ml chloramphenicol , 0.3 mL of antifoam (Sigma), 9 mls of 0.83 M MgSO4 and trace elements in a 1L bottle and cultured using the LEX system to an OD600 of 5, cooled to 15°C, and induced with 0.5 mM isopropyl-1-thio-D-galactopyranoside (IPTG) overnight at 15degC.

Purification

Procedure

The cleared lysate was loaded onto a column prepakced with 10 g DE52 (Whatman) anion

exchange resin (previously activated with 2.5 M NaCl and equilibrated with Binding Buffer); and subsequently onto a 2mL Ni-NTA (Qiagen) column pre-equilibrated with Binding Buffer at approximately 1 - 1.5 mL/min. The volume of the Ni-NTA resin was pre-determined by the predicted protein yield from test expression analysis. After the lysate was loaded, the DE52 was further washed with 20 mL of Binding Buffer. The Ni-NTA column was then washed with 200 mL of Wash Buffer at 2 - 2.5 mL/min. After washing, the protein was eluted with 15 mL of Elution Buffer. 1 mM TCEP and 1 mM EDTA was added to the eluted PBG-PF11_0147.

The protein sample was then loaded onto a superdex 200 gel filtration column. The eluted protein (in 10 mM Hepes, pH 7.5 and 500 mM NaCl) was concentrated using a 15 ml Amicon Ultra centrifugal filter device (Millipore) with a 10 kDa cutoff. The protein was concentrated to 10 mg/ml using a 15 ml Amicon Ultra centrifugal filter device (Millipore) with a 10 kDa cutoff, then stored at 4 °C.

Extraction

Procedure

The culture was harvested by centrifugation. Pellets from 2 L of culture were resuspended to approximately 40 mL/L of cell culture in Binding Buffer with the addition of protease inhibitors (1 mM benzamidine and 1 mM phenylmethyl sulfonyl fluoride (PMSF)). Resuspended pellets stored at -80 degC were thawed overnight at 4 degC on the day before purification. Prior to mechanical lysis, each pellet from 1 L of culture was pretreated with 0.5 % CHAPS and 500 units of benzonase for 40 minutes at room temperature. Cells were mechanically lysed with a microfluidizer (Microfluidizer Processor, M-110EH) at approximately 18000 psi; and the cell lysate was centrifuged using a Beckman JA-25.50 rotor at ~75000 x g (24000 rpms) for 20 minutes at 10 degC.

Concentration:

Ligand

2 mM ATP, 4 mM MgCl₂, 2 mM TCEP

MassSpec:
Crystallization: Crystallization plate: IC-MAZ3MB:C8-1
buffer: 20% PEG 3350, 0.2 M tri-lithium citrate at 20 degC
Cryo conditions: crystallization condition +25% glycerol

NMR Spectroscopy:

Data Collection:

Data Processing: