

**Entry Clone Source:** Synthetic

**Entry Clone Accession:** GI:21071056

**Vector:** pNIC28-Bsa4 Details [[PDF](#)]; Sequence [[FASTA](#)] or [[GenBank](#)]

**Amplified DNA sequence:**

```
CATATGCACCATCATCATCATCATCATTC  
TTCTGGTAGATCTGGTACCGAGA  
ACCTGTACTTCAATCCATGGCGGAA  
AAACTGAGCCGAATCCGCCAACCT  
GACCAAAAAATGAAAAAAATTGTTG  
ATGCGGTGATTAAATATAAGATAGC  
AGCAGCGGTGTCAGCTGAGCGAAGT  
GTTTATTCTAGCTGCCGAGCCGTAAAG  
AACTGCCGGAATATTGAACTGATT  
CGTAAACCGGTGGATTTAAAAAAAT  
TAAAGAACGTATTGCAATCATAAAT  
ATCGTAGCCTGAATGATCTGGAAAAAA  
GATGTTGATGCTGCTGCCAGAATGC  
GCAGACCTTTAACCTGGAAGGCAGCC  
TGATTATGAAGATAGCATTGTTCTG  
CAGAGCGTGTACCGCGTGCAGCCA  
GAAAATCGAAAAAGAAGATGATTGAC  
AGTAAAGGTGGATACGGATCCGAA
```

**Final protein sequence (Tag sequence in lowercase):**

```
mhhhhhhssgvdlgtenlyfq^smAE  
KLSPNPPNLTKKMKKIVDAVIKYKDS  
SSGRQLSEVFIQLPSRKELPEYYELI  
RKPVDFKKIKERIRNHKYRSLNDEK  
DVMLLCQNAQTFNLEGSLIYEDSIVL  
QSVFTSVRQKIEKEDD
```

^ TEV cleavage site

**Tags and additions:** Cleavable N-terminal His6 tag.

**Host:** BL21 (DE3)R3-pRARE2 (Phage resistant strain)

**Growth medium, induction protocol:** 10 ml from a 50 ml overnight culture containing 50  $\mu$ g/ml kanamycin and 34  $\mu$ g/ml chloramphenicol were used to inoculate each of two 1 liter cultures of TB containing 50  $\mu$ g/ml kanamycin and 34  $\mu$ g/ml chloramphenicol. Cultures were grown at 37°C until the OD<sub>600</sub> reached ~2.5 then the temperature was adjusted to 18°C.

Expression was induced overnight using 0.1 mM IPTG at an OD<sub>600</sub> of 3.0. The cells were collected by centrifugation and the pellet re-suspended in binding buffer and frozen.

**Binding buffer:** 50 mM HEPES pH 7.5; 500 mM NaCl; 10 mM imidazole, 5% glycerol

**Extraction buffer, extraction method:** Frozen pellets were thawed and fresh 0.5 mM TCEP, 1 mM PMSF added to the lysate. Cells were lysed using sonication. DNA was precipitated with 0.15% PEI. The lysate was centrifuged at 17,000 rpm for 60 minutes and the supernatant collected for purification.

**Column 1:** Ni-affinity. Ni-sepharose (Amersham), 5 ml of 50% slurry in 1.5 x 10 cm column, washed with binding buffer

**Column 1 Buffers:**

**Binding buffer:** 50 mM HEPES pH 7.5, 500 mM NaCl, 5 mM imidazole, 5% glycerol

**Wash buffer:** 50 mM HEPES pH 7.5, 500 mM NaCl, 30 mM Imidazole, 5% glycerol  
**Elution buffer:** 50 mM HEPES pH 7.5, 500 mM NaCl, 5% glycerol, 50 to 250 mM Imidazole (step elution).

**Column 1 Procedure:** The supernatant was loaded by gravity flow on the Ni-sepharose column. The column was washed first with 30 ml of binding buffer then with 30 ml of wash buffer at gravity flow. The protein was eluted by gravity flow by applying 5ml portions of elution buffer with increasing concentration of imidazole (50 mM, 100 mM, 150 mM, 200 and 250 mM); fractions were collected until essentially all protein was eluted

**Enzymatic treatment:** The N-terminal His tag was cleaved by treatment with TEV protease, overnight.

**Column 2:** Size Exclusion Chromatography; Superdex S75 16/60 HiLoad

**Column 2 Buffers:**

**Buffer:** 10 mM HEPES, pH 7.5; 500 mM NaCl, 5% glycerol

**Column 2 Procedure:** The protein was concentrated and applied to an S75 16/60 HiLoad gel filtration column equilibrated in 10 mM HEPES, pH 7.5; 500mM NaCl, 5% glycerol using an AKTAexpress system.

**Protein concentration:** The protein was concentrated to 11 mg/ml using an Amicon 3 kDa cut-off concentrator.

**Mass spectrometry characterization:**

**Measured mass:** 14535.8 Da

LC- ESI -MS TOF gave a measured mass of 14535.8 Da for the cleaved protein as predicted from the sequence.

**Crystallisation:** Protein buffer was exchanged to 10 mM HEPES pH7.5 and 300 mM NaCl. Crystals were grown at 4°C in 300 nl sitting drops from a 1:1 ratio of protein (11 mg/ml with 5 mM NMP) to reservoir solution containing 0.1M Tris pH 8.5 and 28% MMW PEG-Smear

**Data collection:**

**Resolution:** 1.54Å

Crystals were cryo-protected using the well solution supplemented by 20% ethylene glycol and flash frozen in liquid nitrogen. Data were collected in-house on a Rigaku FRE-Superbright equipped with an RAXIS-IV detector at a wavelength of 1.542 Å.

**Phasing:** The structure was solved by molecular replacement using an ensemble of known bromodomain structures as a starting model